

	Prevents	Birth	6 weeks	10 weeks	14 weeks	6 months	9 months	9-12 months	12 months	15 months	15-18 months	16-18 months	18 months	2 years	4-6 years	9-14 years	10-12 years
BCG	TB	1 <sup>st</sup> Dose															
OPV (Oral vaccine)	Polio	1 <sup>st</sup> Dose															
Hep-B	Hepatitis B	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose			3 <sup>rd</sup> Dose											
DTwP	Diphtheria, Tetanus & Pertussis	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose	3 <sup>rd</sup> Dose								Booster Dose 1			Booster Dose 2		
IPV (Injectable vaccine)	Polio	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose	3 <sup>rd</sup> Dose								Booster Dose					
Hib	Infections caused by bacteria: Haemophilus influenzae type b	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose	3 <sup>rd</sup> Dose								Booster Dose					
Rotavirus	Diarrhoea	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose	3 <sup>rd</sup> Dose													
PCV	Pneumonia	1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose	3 <sup>rd</sup> Dose						Booster Dose							
OPV	Polio					1 <sup>st</sup> Dose	2 <sup>nd</sup> Dose										3 <sup>rd</sup> Dose
MMR	Measles, Mumps & Rubella						1 <sup>st</sup> Dose			2 <sup>nd</sup> Dose							3 <sup>rd</sup> Dose
Typhoid Conjugate Vaccine	Typhoid						1 <sup>st</sup> Dose							Booster Dose			
Hep - A*	Hepatitis A							1 <sup>st</sup> Dose (if LAV or inactivated vaccine)			2 <sup>nd</sup> Dose (if inactivated vaccine)						
Varicella	Chicken pox									1 <sup>st</sup> Dose							2 <sup>nd</sup> Dose*
Tdap/Td	Diphtheria, Tetanus & Pertussis																1 <sup>st</sup> Dose
HPV (for girls only)	Cervical cancer																1 <sup>st</sup> Dose (2nd dose 6 months apart)

\*Hep A - Only one dose if LAV (Live Attenuated Vaccine) and 2 doses if inactivated vaccine

\*\*2nd dose of Varicella vaccine can be given anytime after 3 months from 1<sup>st</sup> dose